

## Bibliography for MS I Feeding Behavior Lecture, 2/19/08

- Bjursell, M., Gerdin, A. K., Lelliott, C. J., Egecioglu, E., Elmgren, A., Tornell, J., et al. (2008). Acutely reduced locomotor activity is a major contributor to Western diet-induced obesity in mice. *Am J Physiol Endocrinol Metab*, 294(2), E251-260.
- CDC. (2008). Obesity and Overweight: Trends: U.S. Obesity Trends 1985-2006 | DNPAO | CDC. Retrieved February 19, 2008, from <http://www.cdc.gov/nccdphp/dnpa/obesity/trend/maps/>
- Cummings, D. E., Weigle, D. S., Frayo, R. S., Breen, P. A., Ma, M. K., Dellinger, E. P., et al. (2002). Plasma ghrelin levels after diet-induced weight loss or gastric bypass surgery. *N Engl J Med*, 346(21), 1623-1630.
- de Castro, J. M., Bellisle, F., Dalix, A. M., & Pearcey, S. M. (2000). Palatability and intake relationships in free-living humans. characterization and independence of influence in North Americans. *Physiol Behav*, 70(3-4), 343-350.
- de Castro, J. M. (2000). Eating behavior: lessons from the real world of humans. *Nutrition*, 16(10), 800-813.
- Elliott, S. (2007, 12/6/2007). Straight A's, With a Burger as a Prize. *New York Times*.
- Elliott, S. S., Keim, N. L., Stern, J. S., Teff, K., & Havel, P. J. (2002). Fructose, weight gain, and the insulin resistance syndrome. *Am J Clin Nutr*, 76(5), 911-922.
- Levine, A. S., Weldon, D. T., Grace, M., Cleary, J. P., & Billington, C. J. (1995). Naloxone blocks that portion of feeding driven by sweet taste in food-restricted rats. *Am J Physiol*, 268(1 Pt 2), R248-252.
- Moran, T. H., & Kinzig, K. P. (2004). Gastrointestinal satiety signals II. Cholecystokinin. *Am J Physiol Gastrointest Liver Physiol*, 286(2), G183-188.
- Pollan, M. (2007, January 28, 2007). Unhappy Meals. *New York Times*.
- Rolls, B. J., Roe, L. S., & Meengs, J. S. (2007). The effect of large portion sizes on energy intake is sustained for 11 days. *Obesity (Silver Spring)*, 15(6), 1535-1543.
- Rolls, B. J., Roe, L. S., & Meengs, J. S. (2006). Larger portion sizes lead to a sustained increase in energy intake over 2 days. *J Am Diet Assoc*, 106(4), 543-549.
- Saper, C. B., Chou, T. C., & Elmquist, J. K. (2002). The need to feed: homeostatic and hedonic control of eating. *Neuron*, 36(2), 199-211.
- Smith, G. P. (2000). The controls of eating: a shift from nutritional homeostasis to behavioral neuroscience. *Nutrition*, 16(10), 814-820.
- Spiegel, T. A., Kaplan, J. M., Tomassini, A., & Stellar, E. (1993). Bite size, ingestion rate, and meal size in lean and obese women. *Appetite*, 21(2), 131-145.
- Teff, K. L., Elliott, S. S., Tschop, M., Kieffer, T. J., Rader, D., Heiman, M., et al. (2004). Dietary fructose reduces circulating insulin and leptin, attenuates postprandial suppression of ghrelin, and increases triglycerides in women. *J Clin Endocrinol Metab*, 89(6), 2963-2972.